

Lassa fever Situation Report

Epi Week 52: 27 December – 2 January, 2021

Key Points

Table 1: Summary of current week (52), cumulative from Epi week 01–52, 2021 and comparison with previous year (2020)

Reporting Period	Suspected cases	Confirmed cases	Probable cases	Deaths (Confirmed cases)	Case Fatality Ratio (CFR)	States and LGAs affected (Confirmed cases)
Current week (week 52)	174	28	1	3	10.7%	State(s): 6 LGA(s): 12
2021 Cumulative (week 1-52)	4654	510	6	102	20.0%	State(s): 17 LGA(s): 68
2020 Cumulative (week 1-52)	6732	1181	14	244	20.7%	State(s): 27 LGA(s): 131

Highlights

- In week 52, the number of new confirmed cases decreased from 29 in week 51, 2021 to 28 cases. These were reported from Ondo, Edo, Bauchi, Kaduna and Taraba States and the FCT (Table 3)
- Cumulatively from week 1 to week 52, 2021, 102 deaths have been reported with a case fatality rate (CFR) of 20.0% which is lower than the CFR for the same period in 2020 (20.7%)
- In total for 2021, 17 States have recorded at least one confirmed case across 68 Local Government Areas (Figure 2 and 3)
- Of all confirmed cases, 84% are from Edo (42%), Ondo (34%) and Bauchi (8%) States.
- The predominant age-group affected is 21-30 years (Range: <1 to 70 years, Median Age: 29 years). The male to female ratio for confirmed cases is 1:0.9 (Figure 4)
- The number of suspected cases has decreased compared to that reported for the same period in 2020
- No new Healthcare workers affected in the reporting week 52
- National Lassa fever multi-partner, multi-sectoral Technical Working Group (TWG) continues to coordinate the response activities at all levels




Figure 1. Epidemic curve of confirmed Lassa fever cases epidemiological week 52, 2021




Figure 2. Confirmed Lassa fever cases by States in Nigeria, week 52, 2021




Figure 3. Confirmed Lassa fever rate per 100,000 population for LGAs in Nigeria, week 52, 2021

Table 2: Key indicators for current week 2021 and trend compared to previous week, Nigeria

Indicator	Number for current week	Trend from previous week	Cumulative number for 2020
Probable cases	1	↔	6
Health Care Worker affected	0	↔ ↔	10
Cases undergoing treatment in Treatment centres	28	↔	482
Contact tracing			
Cumulative contact listed	14	↔	3466
Contacts under follow up	434	↔	434
Contacts completed follow up	67	↔	3006
Symptomatic contacts	0	↔	40
Positive contacts	0	↔	19
Contacts lost to follow up	0	↔ ↔	7

Key

- ↑ Increase
- ↓ Decrease
- ↔ No difference

Table 3. Weekly and Cumulative number of suspected and confirmed cases for 2021

States	Current week: (Week 52)					Cumulative (Week 1 - 52)					
	Cases				Deaths (Confirmed Cases)	Cases				Deaths (Confirmed Cases)	
	Suspected	Confirmed	Trend	Probable		HCW	Suspected	Confirmed	Probable		HCW
1 Edo	74	9	▲			2725	212			15	
2 Ondo	52	12	▲			1006	175	1	6	47	
3 Bauchi	20	4	▼			164	39	1		12	
4 Taraba	1	1	▲			72	22	1	1	12	
5 Ebonyi						143	18			9	
6 Plateau	1		▼			42	9				
7 Benue	3					34	8				
8 Kaduna	2	1	▲	1		53	8	1		4	
9 Enugu						14	5	1	1	1	
10 Nasarawa	12					79	3	1	2	2	
11 Kogi	1					17	3				
12 FCT	4	1	▲			59	3				
13 Cross River						4	1				
14 Imo						9	1				
15 Anambra						10	1				
16 Delta						50	1				
17 Abia						11	1				
18 Plateau						1					
19 Sokoto						3					
20 Kwara						4					
21 Bayelsa						3					
22 Katsina						2					
23 Niger						1					
24 Ogun						3					
25 Oyo						4					
26 Rivers						10					
27 Zamfara						1					
28 Jigawa	1					12					
29 Adamawa						8					
30 Gombe						22					
31 Lagos						16					
32 Kano						24					
33 Ekiti						3					
34 Yobe						4					
35 Kebbi						2					
36 Borno	2					15					
37 Osun						3					
Total	173	28	▼	1	0	3	4633	510	6	10	102

Key	
▼	Decrease
▲	Increase




Figure 4. Age and sex pyramid showing number of confirmed Lassa fever cases for 2021




Figure 5: Number of confirmed cases with case fatality rate (CFR) by state week 52 2021




Figure 6: Trend of confirmed cases by epidemiological week, 2017– 2021, Nigeria

Response activities

- Conducted 2021 Lassa fever high burden States preparedness/response engagement meeting
- Lassa fever alert letters sent to States
- The National Emergency Operations Centre alert mode activated 2021 for effective multi-sectoral, multi-disciplinary coordination of Lassa fever response
- State Public Health Emergency Operations Centre activated in affected States
- The five Lassa fever molecular laboratories in the NCDC network are working full capacity to ensure that all samples are tested and results provided within the shortest turnaround time
- Confirmed cases are treated at identified treatment centres across the states
- Dissemination of reviewed case management and safe burial practices guidelines
- Risk communications and community engagement activities have been scaled up across states using television, radio, print, social media and other strategies
- Implementation of Lassa fever Environmental response campaign in high burden states by Federal Ministry of Environment
- Deployment of National Rapid Respond Teams to three States to support Lassa fever Response

Notes on this report

Data Source

Information for this disease was case based data retrieved from the National Lassa fever Emergency Operations Centre.

Case definitions

- **Suspected case:** any individual presenting with one or more of the following: malaise, fever, headache, sore throat, cough, nausea, vomiting, diarrhoea, myalgia, chest pain, hearing loss and either a. History of contact with excreta or urine of rodents b. History of contact with a probable or confirmed Lassa fever case within a period of 21 days of onset of symptoms OR Any person with inexplicable bleeding/hemorrhagia.
- **Confirmed case:** any suspected case with laboratory confirmation (positive IgM antibody, PCR or virus isolation)
- **Probable case:** any suspected case (see definition above) who died or absconded without collection of specimen for laboratory testing
- **Contact:** Anyone who has been exposed to an infected person, or to an infected person's secretions, excretions, or tissues within three weeks of last contact with a confirmed or probable case of Lassa fever

Calculations

- Case Fatality Rate (CFR) for this disease is reported for confirmed cases only