

www.ncdc.gov.ng



PLOT 800 EBITU UKIWE STREET, JABI ABUJA, NIGERIA. TOLL FREE CALL: 0800-970000-10. E: info@ncdc.gov.ng

SITUATION REPOR

Nigeria Centre For Disease Control (NCDC)

@NCDCgov

TITLE:	2019 LASSA FEVER OUTBREAK SITUATION REPORT
SERIAL NUMBER:	20
EPI-WEEK:	20
DATE:	19 th May 2019

HIGHLIGHTS

- In the reporting Week 20 (13th 19th May, 2019) six new confirmedⁱ cases were reported from three states Edo(2), Ebonyi(2) and Ondo(2) states with one new death from Edo state
- From 1st January to 19th May, 2019, a total of 2504 **suspectedⁱ cases** have been reported from 21 states. Of these, **575 were confirmed positive**, 15 probable and 1914 <u>negative</u> (not a case)-*Table 1*
- Since the onset of the 2019 outbreak, there have been **129** deaths in confirmed cases. Case fatality ratio in confirmed cases is 22.4%
- Twenty-one (21) States (Edo, Ondo, Bauchi, Nasarawa, Ebonyi, Plateau, Taraba, FCT, Adamawa, Gombe, Kaduna, Kwara, Benue, Rivers, Kogi, Enugu, Imo, Delta, Oyo, Kebbi and Cross River) have recorded at least one confirmed case across 82 Local Government Areas *Figure 1*
- 93% of all confirmed cases are from Edo(36%), Ondo(29%), Ebonyi(8%), Bauchi(7%), Taraba(7%) and Plateau(6%) states- *Figure 1*
- Predominant age-group affected is 21-40 years (Range: >1 month to 89 years, Median Age: 32 years) *Figure 6*
- The male to female ratio for confirmed cases is 1.2:1 Figure 6
- In the reporting week 20, no new health care worker was affected. A total of eighteen health care workers have been infected since the onset of the outbreak in eight States Edo (7), Ondo (3), Ebonyi (2), Enugu (1), Rivers (1), Bauchi (1) Benue (1), Plateau (1) and Kebbi(1) with two deaths in Enugu and Edo States
- Five (5) patients are currently being managed at various treatment centres across the country: Irrua Specialist Teaching Hospital(ISTH) treatment Centre (1), Federal Teaching Hospital Abakaliki (2), Federal Medical Centre, Jalingo (1) and Kebbi (1)
- A total of **7446** contacts have been identified from 20 States. Of these 578(7.8%) are currently being followed up, 6800(91.3%) have completed 21 days follow up, while 8(0.1%) were lost to follow up. 119(1.6%) symptomatic contacts have been identified, of which **60(**0.8%) have tested positive
- Multi sectoral one health national rapid response teams (NCDC, NFELTP and Federal Ministry of Environment) deployed to Kebbi state
- National Lassa fever multi-partner, multi-sectoral Technical Working Group (TWG) continues to coordinate the response activities at all levels

Giz Sectore

BNITM

UK-PHRST AHRICA.CDC

REFEARD & Word Health TREETER COC CONTRACT DURING STATUS




Figure 1. Distribution of Confirmed Lassa Fever cases in Nigeria as at 19th May, 2019




Figure 2. LGA attack rate of confirmed Lassa fever cases in Nigeria as at 19th May, 2019






Figure 3. Epicurve of Lassa fever Confirmed (575) Cases in Nigeria - week 01-20, 2019




Figure 4. Weekly trends of Lassa fever Confirmed Cases in Nigeria, 2016-2019/week 20






Figure 5. Confirmed Lassa fever cases in Nigeria with state specific Case Fatality Rates (CFR) as at 19th May, 2019




Figure 6. Age-Sex distribution of Confirmed Lassa fever cases in Nigeria as at 19th May, 2019

<u>Disclaimer –</u> The information contained in this document is confidential, privileged and only for the intended recipient and may not be used, published or redistributed to the public. A redacted version is available on <u>http://ncdc.gov.ng/diseases/sitreps</u>



ⁱSuspected case describes any individual presenting with one or more of the following: malaise, fever, headache, sore throat, cough, nausea, vomiting, diarrhoea, myalgia, chest pain, hearing loss and either a. History of contact with excreta or urine of rodents b. History of contact with a probable or confirmed Lassa fever case within a period of 21 days of onset of symptoms OR Any person with inexplicable bleeding/hemorrhagia.

ⁱⁱAny suspected case with laboratory confirmation (positive IgM antibody, PCR or virus isolation)

ⁱⁱⁱAny suspected case (see definition above) who died without collection of specimen for laboratory testing

^{iv} "Active" means where there has been at least one confirmed case, and contacts within 21 days post exposure