



Nigeria Centre for Disease Control

Protecting the health of Nigerians

Epi Week: 36 2021

www.ncdc.gov.ng

PLOT 801 EBITU UKIWE STREET, JABI ABUJA, NIGERIA. **TOLL FREE CALL**: 6232. **Email**: info@ncdc.gov.ng

@NCDCgov

Lassa fever Situation Report

Epi Week 36: 6 – 12 September 2021


Key Points

Table 1: Summary of current week (36), cumulative from Epi week 01–36, 2021 and comparison with previous year (2020)

Reporting Period	Suspected cases	Confirmed cases	Probable cases	Deaths (Confirmed cases)	Case Fatality Ratio (CFR)	States and LGAs affected (Confirmed cases)		
Current week (week 36)	60	4	0	3	75.0%	State(s): 2 LGA(s): 4		
2021 Cumulative (week 1-36)	3006	369	3	76	20.6%	State(s): 14 LGA(s): 62		
2020 Cumulative (week 1-36)	5647	1078	14	225	20.9%	State(s): 27 LGA(s): 129		

Highlights

- In week 36, the number of new confirmed cases increased from 0 in week 35, 2021 to 4 cases (Table 3)
- Cumulatively from week 1 to week 36, 2021, 76 deaths have been reported with a case fatality rate (CFR) of 20.6% which is lower than the CFR for the same period in 2020 (20.9%)
- In total for 2021, 14 States have recorded at least one confirmed case across 62 Local Government Areas (Figure 2 and 3)
- Of all confirmed cases, 84% are from Edo (45%), Ondo (34%) and Taraba (5%) States.
- The predominant age-group affected is 21-30 years (Range: <1 to 70 years, Median Age: 29 years). The male to female ratio for confirmed cases is 1:0.9 (Figure 4)
- The number of suspected cases has decreased compared to that reported for the same period in 2020
- No new Healthcare worker affected in the reporting week 36
- National Lassa fever multi-partner, multi-sectoral Technical Working Group (TWG) continues to coordinate the response activities at all levels



Epi Week: 36 2021

Figure 1. Epidemic curve of confirmed Lassa fever cases epidemiological week 36, 2021




Figure 2. Confirmed Lassa fever cases by States in Nigeria, week 36, 2021




Figure 3. Confirmed Lassa fever rate per 100,000 population for LGAs in Nigeria, week 36, 2021

Table 2: Key indicators for current week 2021 and trend compared to previous week, Nigeria

Indicator	Number for current week	Trend from previous week	Cumulative number for 2020	
Probable cases	0	←→	3	
Health Care Worker affected	0	←→	6	
Cases undergoing treatment in Treatment centres	4	↑	343	
Contact tracing			•	
Cumulative contact listed	0	←→	2108	
Contacts under follow up	140	1	140	
Contacts completed follow up	0	1	1953	
Symptomatic contacts	0	←→	8	
Positive contacts	0	←→	8	
Contacts lost to follow up	0	←→	7	


Key Increase Decrease No difference

Table 3. Weekly and Cumulative number of suspected and confirmed cases for 2021

Epi Week: 36 2021

			Cui	k 36)	Cumulative (Week 1 - 36)							
	Chahaa	Cases					Deaths		Cases			Deaths
	States	Suspected	Confirmed	Trend	Probable	HCW ³	(Confirmed Cases)	Suspected	Confirmed	Probable	HCW ³	(Confirmed Cases)
1	Edo	46						1807	165			14
2	Ondo	11	(3 🔺			3	674	124	1	4	4:
3	Taraba	1	:	1				66	20	1	1	1:
4	Ebonyi							114	17			
5	Bauchi							65	10			
6	Plateau							26	8			
7	Benue							23	6			
8	Kaduna							42	6			
9	Enugu							14	5	1	1	
10	Nasarawa							10	3			
11	Kogi							10	2			
	Cross River							3	1			
	Delta							36	1			
14	Abia							8	1			
	Kwara							2				
	Bayelsa							2				
	Katsina							1				
	Niger							1				
	Ogun							3				
	Oyo							2				
	Rivers							9				
	Zamfara							1				
	Jigawa							8				
	Adamawa							7				
	Gombe	1						9				
	Lagos							13				
	Kano							9				
	Ekiti							3				
	Yobe							3				
	FCT							8				
	Imo	1						7				
	Kebbi	1						2				
	Borno							8				
	Anambra							5				
	Osun							2				
35	Journ							2				
	Total	60	,	4 🔺	0	0	3	3003	369	3	6	7(





Epi Week: 36 2021

Figure 4. Age and sex pyramid showing number of confirmed Lassa fever cases for 2021




Figure 5: Number of confirmed cases with case fatality rate (CFR) by state week 36 2021




Figure 6: Trend of confirmed cases by epidemiological week, 2017-2021, Nigeria

Response activities

Conducted 2021 Lassa fever high burden States preparedness/response engagement meeting

Epi Week: 36 2021

- Lassa fever alert letters sent to States
- The National Emergency Operations Centre alert mode activated 2021 for effective multisectoral, multi-disciplinary coordination of Lassa fever response
- State Public Health Emergency Operations Centre activated in affected States
- The five Lassa fever molecular laboratories in the NCDC network are working full capacity to ensure that all samples are tested and results provided within the shortest turnaround time
- · Confirmed cases are treated at identified treatment centres across the states
- Dissemination of reviewed case management and safe burial practices guidelines
- Risk communications and community engagement activities have been scaled up across states using television, radio, print, social media and other strategies
- Implementation of Lassa fever Environmental response campaign in high burden states by Federal Ministry of Environment
- Deployment of National Rapid Respond Teams to three States to support Lassa fever Response

Notes on this report

Data Source

Information for this disease was case based data retrieved from the National Lassa fever Emergency Operations Centre.

Case definitions

- Suspected case: any individual presenting with one or more of the following: malaise, fever, headache, sore throat, cough, nausea, vomiting, diarrhoea, myalgia, chest pain, hearing loss and either a. History of contact with excreta or urine of rodents b. History of contact with a probable or confirmed Lassa fever case within a period of 21 days of onset of symptoms OR Any person with inexplicable bleeding/hemorrhagia.
- **Confirmed case**: any suspected case with laboratory confirmation (positive IgM antibody, PCR or virus isolation)
- Probable case: any suspected case (see definition above) who died or absconded without collection of specimen for laboratory testing
- Contact: Anyone who has been exposed to an infected person, or to an infected person's secretions, excretions, or tissues within three weeks of last contact with a confirmed or probable case of Lassa fever

Calculations

Case Fatality Rate (CFR) for this disease is reported for confirmed cases only