

# SITUATION REPORT

Nigeria Centre For Disease Control and Prevention (NCDC)

NCDC.GOV.NG

PLOT 800 EBITU UKIWE STREET, JABI ABUJA, NIGERIA TOLL FREE CALL: 6232 E:info@ncdc.gov.ng

@NCDCgov 🖪 💆 🖸

TITLE:	UPDATE ON MONKEYPOX (MPX) IN NIGERIA
SERIAL NUMBER:	23
EPI-WEEK:	39
DATE:	October 2, 2022

#### Table 1 – Key Indicators


Key Indicators	Number		
Total confirmed cases in Epi Week 39, 2022			
Total suspected cases from January 1 <sup>st</sup> to October 2 <sup>nd</sup> , 2022 (Epi week 1 to 39)	1180		
Total confirmed cases from January 1 <sup>st</sup> to October 2 <sup>nd</sup> , 2022 (Epi week 1 to 39)	481		
Total deaths from January 1 <sup>st</sup> to October 2 <sup>nd</sup> , 2022 (Epi week 1 to 39)	7		
Total deaths Sept 2017 - October 2 <sup>nd</sup> , 2022	15		
Total confirmed cases in 2017	88		
Total confirmed cases in 2018	49		
Total confirmed cases in 2019	47		
Total confirmed cases in 2020	8		
Total confirmed cases in 2021	34		
Grand total confirmed cases (Sept 2017 – October 2 <sup>nd</sup> , 2022)	707		
Grand total suspected cases (Sept 2017 – October 2 <sup>nd</sup> , 2022)	1692		

- Seventy-two (72) new suspected cases reported in Epi week 39, 2022 (26<sup>th</sup> September to 2<sup>nd</sup> October 2022) from Fifteen (15) states and FCT Lagos (28), Benue (8), Ogun (6), Ondo (5), FCT (4), Kaduna (4), Katsina (3), Borno (2), Delta (2), Ekiti (2), Kebbi (2), Osun (2), Kano (1), Oyo (1), Plateau (1) and Rivers (1).
- Twenty-five (25) new confirmed cases were recorded in Epi week 39 from Nine (9) states and FCT Lagos (7), Benue (5), FCT (3), Borno (2), Delta (2), Kaduna (2), Kebbi (1), Osun (1), Oyo (1) and Rivers (1).
- From 1<sup>st</sup> January to 2<sup>nd</sup> September 2022, Nigeria has recorded 1180 suspected cases with 481 confirmed cases (320 male, 161 female) from thirty-one (31) states and FCT Lagos (123), Abia (32), Ondo (27), Bayelsa (26), Rivers (26), Imo (22), Delta (22), Edo (19), Ogun (19), Adamawa (15), FCT (15), Anambra (15), Nasarawa (12), Benue (10), Plateau (9), Oyo (9), Kwara (8), Borno (8), Kaduna (8), Taraba (7), Katsina (7), Akwa Ibom (7), Cross River (7), Ebonyi (6), Kano (5), Gombe (5), Kogi (4), Osun (4), Kebbi (2), Niger (1) and Bauchi (1).
- Seven (7) associated deaths were recorded from 7 states in 2022 Delta (1), Lagos (1), Ondo (1) and Akwa Ibom (1), Kogi (1), Taraba (1) and Imo (1). **CFR 1.5%**





- Overall, since the re-emergence of monkeypox in September 2017, 1692 suspected cases have been reported from 35 states and FCT in the country. Of these 1620 suspected cases, there have been 707 (41.7%) confirmed (473 male, 234 female) from 32 states and FCT Lagos (153), Rivers (78), Bayelsa (69), Delta (51), Abia (35), Imo (30), Edo (29), Ondo (27), FCT (21), Cross River (21), Ogun (20), Anambra (17), Adamawa (15), Oyo (15), Nasarawa (14), Akwa Ibom (14), Plateau (12), Benue (12), Kwara (8), Borno (8), Kaduna (8), Taraba (7), Katsina (7), Ebonyi (7), Kano (5), Gombe (5), Enugu (4), Kogi (4), Osun (4), Ekiti (2), Niger (2), Kebbi (2)and Bauchi (1).
- Fifteen (15) deaths have been recorded since September 2017 (CFR= 2.1%) in eleven states Lagos (3), Edo (2), Imo (2), Cross River (1), FCT (1), Rivers (1), Ondo (1) Delta (1), Akwa Ibom (1), Taraba (1) and Kogi (1).



PARENET APRICACIA (TARACIA)

Figure 1: Epidemic Curve of Suspected & Confirmed MPX Cases Jan. 2022 till date







Figure 2: Bar chart Showing confirmed monkeypox cases by state from January  $1^{st} - 2^{nd}$  October 2022





*Figure 3: Bar chart Showing confirmed monkeypox cases by state, September 2017 – 2<sup>nd</sup> October 2022* 


Ceffeelth TAFENET AND CONTRACTOR CONTRACTOR




Table 2: Nigeria confirmed	l monkevpox cases b	ov state. September	2017 – 2 <sup>nd</sup> October 2022
Tuble 2. Nigeria conjinnea	monneypox cuses b	y state, september	2017 2 0000007 2022

State	2017	2018	2019	2020	2021	2022	Total
Lagos	4	1	15	4	6	123	153
Rivers	25	14	7	1	5	26	78
Bayelsa	19	11	7	0	6	26	69
Delta	3	6	10	1	9	22	51
Abia	1	2	0	0	0	32	35
Imo	5	2	1	0	0	22	30
Edo	4	1	1	0	4	19	29
Ondo	0	0	0	0	0	27	27
FCT	5	0	0	0	1	15	21
Cross River	9	3	1	0	1	7	21
Ogun	0	0	0	0	1	19	20
Anambra	0	1	1	0	0	15	17
Adamawa	0	0	0	0	0	15	15
Оуо	1	3	2	0	0	9	15
Nasarawa	1	1	0	0	0	12	14
Akwa Ibom	6	0	1	0	0	7	14
Benue	2	0	0	0	0	10	12
Plateau	0	2	0	1	0	9	12
Borno	0	0	0	0	0	8	8
Kaduna	0	0	0	0	0	8	8
Kwara	0	0	0	0	0	8	8
Taraba	0	0	0	0	0	7	7
Katsina	0	0	0	0	0	7	7
Ebonyi	0	0	0	1	0	6	7
Kano	0	0	0	0	0	5	5
Gombe	0	0	0	0	0	5	5
Enugu	1	2	1	0	0	0	4
Kogi	0	0	0	0	0	4	4
Osun	0	0	0	0	0	4	4
Niger	0	0	0	0	1	1	2
Ekiti	2	0	0	0	0	0	2
Kebbi	0	0	0	0	0	2	2
Bauchi	0	0	0	0	0	1	1
Grand Total	88	49	47	8	34	481	707






*Figure 4:* Map of Nigeria Showing States with Confirmed MPX Cases from January 2022 till date (30 states and FCT)



*Figure 5:* Map of Nigeria Showing States with Confirmed MPX Cases from September 2017 till date (32 states and FCT)

World Health Organization






*Figure 6:* Nigeria confirmed Monkeypox cases by the year of incidence- September 2017 to 2<sup>nd</sup> October 2022

able 5. Age distribution of confirmed Monkeypox cases September 2017 – 2 October 2022							
Age Group	2017	2018	2019	2020	2021	2022	Total
0-10 Years	7	5	1	0	1	72	86
11-20 Years	12	4	1	0	4	76	97
21-30 Years	34	13	13	4	10	115	189
31- 40 Years	26	17	22	4	13	141	223
41-50 Years	9	10	9	0	5	57	90
51-60 Years	0	0	1	0	1	20	22
Total	88	49	47	8	34	481	707


Table 3: Age distribution of confirmed Monkeypox cases September 2017 – 2<sup>nd</sup> October 2022







*Figure 7:* Age and sex distribution of Nigeria confirmed monkeypox cases January 1<sup>st</sup> – 2<sup>nd</sup> October 2022



*Figure 8:* Age and sex distribution of Nigeria confirmed monkeypox cases September 2017 – 2<sup>nd</sup> October 2022



Pillar	Activities to date	Next steps
Coordination	<ul> <li>Supporting the daily activities of the activated Emergency Operations Centre (EOC)</li> </ul>	<ul> <li>The EOC will continue coordinating ongoing response activities in the country while contributing to the global response.</li> </ul>
Risk communication	<ul> <li>Daily tracking, monitoring and response to rumours and misinformation of Monkeypox in Nigeria</li> <li>Collaboration with partners to develop Monkeypox-specific awareness content</li> </ul>	<ul> <li>Continue monitoring and analysis of Monkeypox infodemics</li> <li>Continue engagement of social media channels with key messages on Monkeypox prevention</li> </ul>
Surveillance	<ul> <li>Providing off-site support to states on case identification, reporting and response</li> </ul>	<ul> <li>Ensure timely uploading of cases to the SORMAS platform</li> <li>To provide off site support to states on conduct of active case searches in facilities and communities</li> <li>Follow-up calls/emails to the state's surveillance team and MPX treatment facilities.</li> </ul>
IPC	<ul> <li>Development of a screening and triage tool on MPX for health facilities</li> <li>Engagement meeting with U.S CDC team on wide range of support to the National IPC program and Orange Network</li> <li>MPX HCW infection surveillance continues across the Orange Network sites and states</li> <li>Formation of a small technical working group on MPX IPC guideline development</li> </ul>	<ul> <li>Training of 25 IPC FPs in two (2) states (Plateau and Benue) from APIN- Orange Network on IPC workplan development and MPX</li> <li>Continuous support to states and facilities on IPC program</li> </ul>
Case management	<ul> <li>Following up with all positive cases at the state level</li> </ul>	<ul> <li>Ensure up-to-date statistics on cases at Treatment Centre and those on home-based care</li> </ul>
POE	Screening at points of entry for persons of Interest	<ul> <li>Continue ongoing screening activities at points of entry</li> </ul>



Laboratory	<ol> <li>All MPX virus sequences are all clade II</li> <li>Off-site support to states on sample collection from sites to testing lab.</li> </ol>	<ul> <li>To shorten turnaround time of samples from point of collection to arrival at NRL</li> </ul>
------------	--	--

## Notes on this report

#### Data Source

Information for this disease was case-based data retrieved from the National Monkeypox Emergency Operations Centre.

## **Case definitions**

## Suspected case

 An acute illness with fever >38.3°C, intense headache, lymphadenopathy, back pain, myalgia, and intense asthenia followed one to three days later by a progressively developing rash often beginning on the face (most dense) and then spreading elsewhere on the body, including soles of feet and palms of the hand

#### Probable case

• A case that meets the clinical case definition is not laboratory-confirmed but has an epidemiological link to a confirmed case

## Confirmed case

• A clinically compatible case that is laboratory confirmed

#### Contact

• Any person who has been in direct or indirect contact with a confirmed case since the onset of symptoms, i.e., contact with skin lesions, oral secretions, urine, faeces, vomitus, blood, sexual contact, sharing a common space (anyone who has been in proximity with or without physical contact with a confirmed case)

Ochealth TAFENET AREA TA

## Calculations

• Case Fatality Rate (CFR) for this disease is reported for confirmed cases only

