



Nigeria Centre for Disease Control

Protecting the health of Nigerians

Epi Week: 4, 2022

www.ncdc.gov.ng

PLOT 801 EBITU UKIWE STREET, JABI ABUJA, NIGERIA. **TOLL FREE CALL**: 6232. **Email**: info@ncdc.gov.ng

@NCDCgov

Lassa fever Situation Report

Epi Week 4: 24 – 30 January, 2022

Key Points

Table 1: Summary of current week (4), cumulative from Epi week 1−4, 2022 and comparison with previous year (2022)

•								
Reporting Period	Suspected cases	Confirmed cases	Probable cases	Deaths (Confirmed cases)	Case Fatality Ratio (CFR)	States and LGAs affected (Confirmed cases)		
Current week (week 4)	229	42	1	6	14.3%	State(s): 5 LGA(s): 16		
2022 Cumulative (week 4)	981	211	2	40	19.0%	State(s): 14 LGA(s): 43		
2021 Cumulative (week 4)	507	54	0	12	22.2%	State(s): 8 LGA(s): 23		

Highlights

- In week 4, the number of new confirmed cases decreased from 74 in week 3, 2022 to 42 cases. These were reported from Ondo, Bauchi, Edo, Enugu and Delta States (Table 3)
- Cumulatively from week 1 to week 4, 2022, 40 deaths have been reported with a case fatality rate (CFR) of 19.0% which is lower than the CFR for the same period in 2021 (22.2%)
- In total for 2022, 14 States have recorded at least one confirmed case across 43 Local Government Areas (Figure 2 and 3)
- Of all confirmed cases, 82% are from Ondo (30%), Edo (27%) and Bauchi (25%) States.
- The predominant age-group affected is 21-30 years (Range: 1 to 80 years, Median Age: 30 years). The male to female ratio for confirmed cases is 1:0.8 (Figure 4)
- The number of suspected cases has increased compared to that reported for the same period in 2021
- No new Healthcare worker affected in the reporting week 4
- National Lassa fever multi-partner, multi-sectoral Emergency Coordination Centre (EOC) activated to coordinate response activities at all levels




Figure 1. Epidemic curve of confirmed Lassa fever cases epidemiological week 4, 2022




Figure 2. Confirmed Lassa fever cases by States in Nigeria, week 4, 2022




Figure 3. Confirmed Lassa fever rate per 100,000 population for LGAs in Nigeria, week 4, 2022

Indicator	Number for current week	Trend from previous week	Cumulative number for 2022		
Probable cases	1	Ø	2		
Health Care Worker affected	0	0 0	4		
Cases undergoing treatment in Treatment centres	42	Ø	233		
C ontact tracing	•				
C umulative contact listed	339	Ø	968		
Contacts under follow up	617	N	617		
Contacts completed follow up	81	Ø	316		
Symptomatic contacts	0	Ø Ø	44		
Positive contacts	0	0 0	35		
Contacts lost to follow up	0	N N	0		

Table 2: Key indicators for current week 2021 and trend compared to previous week, Nigeria

Increase
Decrease
No difference

Table 3. Weekly and Cumulative number of suspected and confirmed cases for 2021

		Current week: (Week 4)					Cumul	ative (W eek	1 - 4	1)
States Suspecte		Cases					Cases			Deaths
	Suspected	Confirmed	Trend	Probable HCW *	(Confirmed Cases)	Suspected	Confirmed	Probable H	CW *	(Confirmed Cases)
1 Ondo	44	15	•		2	181	63			1:
2 Edo	63	10	▼			317	57		3	!
3 Bauchi	89	14	•			248	53			
4 Benue			▼	1	2	50	11	1	2	
5 O y o	10		▼			25	5			:
6 Taraba			▼			15	5			
7 Ebonyi	7		V			23	5	1		:
8 Kogi						7	4			
9 Kaduna			▼			19	2			-
10 Enugu	5	2	A		2	9	2			
11 Delta	5	1	A			7	1			
12 Katsina			•			2	1			•
13 Plateau						13	1			
14 Cross River	r					3	1			
15 Borno						2				
16 Anambra						2				
17 Bayelsa						1				
18 Jigawa						1				
19 Kebbi						1				
20 O gu n						3				
21 Kwara	1					3				
22 Lagos	2					11				
23 Gombe						6				
24 FCT						15		,		
25 Nasarawa						11				
26 Rivers	3					4				
Total	229	42	_	1 0	6	979	211	2	5	40






Figure 4. Age and sex pyramid showing number of confirmed Lassa fever cases for 2022




Figure 5: Number of confirmed cases with case fatality rate (CFR) by state week 4, 2022




Figure 6: Trend of confirmed cases by epidemiological week, 2018 - 2022, Nigeria

Response activities

Lassa Fever Situation Report

- Lassa fever alert letters sent to States
- The National Emergency Operations Centre response mode Level 2 activated for effective multisectoral, multi-disciplinary coordination of 2022 Lassa fever outbreak response
- Lassa fever preparedness assessment carried out for 36 States and FCT
- State Public Health Emergency Operations Centre activated in affected States
- The Seven Lassa fever molecular laboratories in the NCDC network are working full capacity to ensure that all samples are tested, and results provided within the shortest turnaround time
- Confirmed cases are treated at identified treatment centres across the states
- Dissemination of reviewed case management and safe burial practices guidelines
- Dissemination of reviewed IPC guideline and health facility IPC advisory
- Risk communications and community engagement activities have been scaled up across states using television, radio, print, social media and other strategies
- Implementation of Lassa fever Environmental response campaign in high burden states by Federal Ministry of Environment
- Distribution of medical response commodities to states and treatment centre
- Deployment of National Rapid Respond Teams (NRRT) deployment to Nasarawa, FCT, Edo, Ondo, Bauchi, Ebonyi, Oyo, Taraba, and Benue

Notes on this report

Data Source

Information for this disease was case based data retrieved from the National Lassa fever Emergency Operations Centre.

Case definitions

- Suspected case: any individual presenting with one or more of the following: malaise, fever, headache, sore throat, cough, nausea, vomiting, diarrhoea, myalgia, chest pain, hearing loss and either a. History of contact with excreta or urine of rodents b. History of contact with a probable or confirmed Lassa fever case within a period of 21 days of onset of symptoms OR Any person with inexplicable bleeding/hemorrhagia.
- Confirmed case: any suspected case with laboratory confirmation (positive IgM antibody, PCR or virus
- Probable case: any suspected case (see definition above) who died or absconded without collection of specimen for laboratory testing
- Contact: Anyone who has been exposed to an infected person, or to an infected person's secretions, excretions, or tissues within three weeks of last contact with a confirmed or probable case of Lassa fever

Calculations

Case Fatality Rate (CFR) for this disease is reported for confirmed cases only

























Epi Week: 4, 2022

